

## Ki67 Ab

[References\(62\)](#) [Images\(45\)](#)

Cat.#: AF0198  
Size: 100ul,200ul,50ul

Concn.: ~1mg/ml  
Source: Rabbit

Mol.Wt.: 358kDa  
Clonality: Polyclonal

Application: WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500, ELISA(peptide)  
1:20000-1:40000

\*The optimal dilutions should be determined by the end user.

Reactivity: Human,Mouse,Rat


Purification: The antiserum was purified by peptide affinity chromatography using  
SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

Immunogen: A synthesized peptide derived from human Ki67, corresponding to a region  
within the internal amino acids.

Uniprot: P46013

Description: KI-67 a protein that may be a marker of proliferating cells, involved in  
chromatin compaction. Its expression is altered in many tumor types  
including osteosarcomas, histiocytomas, prostate, breast and esophageal  
cancers. Mutated in colon, cervical and lung cancers.

Storage: Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02%  
sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from  
date of receipt.




Western blot analysis of extracts from various samples, using Ki67 Ab.


Lane 1: Mouse brain lysates;

Lane 2: Rat spleen lysates;

Lane 3: Rat spleen lysates treated with blocking peptide;



AF0198 at 1/100 staining Human normal tissues adjacent to mammary cancer  
by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen  
retrieval step in citrate buffer was performed. The sample was then blocked  
and incubated with the primary Ab at 4°C overnight. An HRP conjugated anti-  
Rabbit Ab was used as the secondary Ab.



AF0198 staining MCF-7 cells by ICC/IF. Cells were fixed with PFA and permeabilized in 0.1% saponin prior to blocking in 10% serum for 45 minutes at 37°C. The primary Ab was diluted 1/400 and incubated with the sample for 1 hour at 37°C. A Alexa Fluor® 594 conjugated goat polyclonal to rabbit IgG (H+L), diluted 1/600 was used as secondary Ab.

**IMPORTANT:** For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.