

HDAC2 Ab


Affinity Biosciences website:www.affbiotech.com order:order@affbiotech.com

References(1) Images(4)

Cat.#: AF6470 Size: 100ul,200ul,50ul	Concn.: ~1mg/ml Source: Rabbit	Mol.Wt.: 55kDa Clonality: Polyclonal
Application:	WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000	
Reactivity:	*The optimal dilutions should be determined by the end user. Human,Mouse,Rat	
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling Resin (Thermo Fisher Scientific).	
Immunogen:	A synthesized peptide derived from human HDAC2, corresponding to a region within the internal amino acids.	
Uniprot:	Q92769	
Description:	HDAC2 a transcriptional regulator of the histone deacetylase family, subfamily 1. Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation plays a role in epigenetic repression and transcriptional regulation, cell cycle progression and developmental events.	
Storage:	Rabbit IgG in phosphate buffered saline sodium azide and 50% glycerol. Store a date of receipt.	



Western blot analysis of extracts from MCF7, using HDAC2 Ab. Lane 1 was treated with the blocking peptide.



AF6470 staining Hela cells by IF/ICC. The samples were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. Samples were then incubated with primary Ab(AF6470 1:200) and mouse anti-beta tubulin Ab(T0023 1:200) for 1 hour at 37°C. An AlexaFluor594 conjugated goat anti-rabbit IgG(H+L) Ab(Red) and an AlexaFluor488 conjugated goat anti-mouse IgG(H+L) Ab(Green) were used as the secondary Ab. The nuclear counter stain is DAPI(blue).

IMPORTANT: For western blot, incubate membrane with diluted primary Ab



Affinity Biosciences website:www.affbiotech.com order:order@affbiotech.com

in 5% w/v milk , 1X TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.