

ACTH (Adrenocorticotropic Hormone) (Polyclonal)

Rabbit anti-human ACTH (Adrenocorticotrophic Hormone) Polyclonal Antibody

REFERENCES AND PRESENTATIONS¹

 ready-to-use (manual or LabVision AutoStainer)

MAD-005138QD-3 MAD-005138QD-7 MAD-005138QD-12

Ready-to-use (MD-Stainer)²
 MAD-005138QD-3/V
 MAD-005138OD/V

concentrated

MAD-005138Q - 1:50 recommended dilution

COMPOSITION

Anti-human ACTH rabbit polyclonal antibody purified from serum and prepared in 10mM PBS, pH 7.4, with 0.2% BSA and 0.09% sodium azide

INTENDED USE : Immunohistochemistry (IHC) on paraffin embedded tissues. Not tested on frozen tissues or Western-Blotting

CLONE: Polyclonal **Ig ISOTYPE:** Rabbit IgG

IMMUNOGEN: Synthetic peptide derived from N-

terminus of human ACTH.

SPECIES REACTIVITY: In vitro diagnostics in humans.

Not tested in other species

DESCRIPTION AND APPLICATIONS:

Anti-ACTH is a useful marker in classification of pituitary tumors and the study of pituitary disease. It reacts with ACTH-producing cells (corticotrophs). It also may react with other tumors (e.g., some small cell carcinomas of the lung) causing paraneoplastic syndromes by secreting ACTH.

IHC POSITIVE CONTROL: Hypophysis **VISUALIZATION:** Cell cytoplasm

IHC RECOMMENDED PROCEDURE:

 - 4μm thick section should be taken on charged slides; dry overnight at 60°C

sindes; dry overnight at 60°C

- Deparaffinise, rehydrate and HIER (heat induced epitope retrieval) boil tissue in the Pt Module using Vitro S.A CITRATE buffer pH6³ for 20 min at 95°C. Upon completion rinse with 3-5 changes of distilled or deionised water followed
- Endogenous peroxidase block Blocking for 10 minutes at room temperature using peroxidase solution (ref. MAD-021540Q-125)

by cooling at RT for 20 min

- Primary antibody: incubate for 10 minutes [The antibody dilution (when concentrated) and protocol may vary depending on the specimen preparation and specific application. Optimal conditions should be determined by the individual laboratory]
- For detection use Master Polymer Plus Detection System (HRP) (DAB included; ref. MAD-000237QK)
- Counterstaining with haematoxylin and final mounting of the slide

STORAGE AND STABILITY: Stored at 2-8°C. Do not freeze. Once the packaging has been opened it can be stored until the expiration date of the reagent indicated on the label. If the reagent has been stored under other conditions to those indicated in this document, the user must first check its correct performance taking into account the product warranty is no longer valid.

WARNINGS AND PRECAUTIONS:

- 1. Avoid contact of reagents with eyes and mucous membranes. If reagents come into contact with sensitive areas, wash with copious amounts of water.
- 2. This product is harmful if swallowed.
- 3. Consult local or state authorities with regard to recommended method of disposal.
- 4. Avoid microbial contamination of reagents.

SAFETY RECOMMENDATIONS

This product is intended for laboratory professional use only. The product is NOT intended to be used as a drug or for domestic purposes. The current version of the Safety Data Sheet for this product can be downloaded by searching the reference number at www.vitro.bio or can be requested at regulatory@vitro.bio.

³ Ref: MAD-004071R/D



 ϵ



¹ These references are for presentation in vials of Low Density Polyethylene (LDPE) dropper. In case the products are used in automated stainers, a special reference is assigned as follows:

^{- /} L: Cylindrical screw-cap vials (QD-3 / L, QD-7 / L, QD-12 / L).
- / N: Polygonal screw-cap vials (QD-3 / N, QD-7 / N, QD-12 / N).
For different presentations (references / volumes) please contact the supplier.

² For Technical specifications for MD-Stainer, please contact your distributor.



BIBLIOGRAPHY

- 1. Hsu DW; et al. American Journal of Pathology, 1991 Apr, 138(4):897-909.
- 2. Kasper M; et al. Journal fur Hirnforschung,1991, 32(6):725-34

LABEL AND BOX SYMBOLS

Explanation of the symbols of the product label and box:

	Expiration date
Î	Temperature limit
***	Manufacturer
Σ	Sufficient content for <n> assays</n>
REF	Catalog number
LOT	Lot code
[]i	Refer to the instructions of use
IVD	Medical product for <i>in</i> vitro diagnosis.
e-SDS	Material safety data sheet

