Leader in Biomolecular Solutions for Life Science

PSMA6 Rabbit pAb

Catalog No.: A13536



Basic Information

Observed MW 27kDa

Calculated MW 18kDa/25kDa/27kDa

Category Primary antibody

Applications WB,IF/ICC

Cross-Reactivity Human, Mouse, Rat

Background

The proteasome is a multicatalytic proteinase complex with a highly ordered ringshaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a nonlysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the peptidase T1A family, that is a 20S core alpha subunit. Multiple transcript variants encoding several different isoforms have been found for this gene. A pseudogene has been identified on the Y chromosome.

Recommended Dilutions

WB 1:500 - 1:2000 IF/ICC 1:50 - 1:200

Immunogen Information

Gene ID 5687 Swiss Prot P60900

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-246 of human PSMA6 (NP_002782.1).

Synonyms

PSMA6;IOTA;PROS27;p27K

Contact	
---------	--

Product Information

Ð	www.abclonal.com
---	------------------

lsotype IgG Purification Affinity purification

Storage

Source

Rabbit

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Validation Data



Western blot analysis of extracts of various cell lines, using PSMA6 antibody (A13536) at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25ug per lane.

Blocking buffer: 3% nonfat dry milk in TBST.



Immunofluorescence analysis of MCF-7 cells using PSMA6 antibody (A13536). Blue: DAPI for nuclear staining.