

ABCG1 Rabbit pAb

Catalog No.: A4328 **4 Publications**

Basic Information

Observed MW

100kDa

Calculated MW

71kDa/73kDa/74kDa/75kDa/86kDa

Category

Primary antibody

Applications

WB, IHC, IF/ICC

Cross-Reactivity

Human, Mouse, Rat

Background

The protein encoded by this gene is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the White subfamily. It is involved in macrophage cholesterol and phospholipids transport, and may regulate cellular lipid homeostasis in other cell types. Six alternative splice variants have been identified.

Recommended Dilutions

WB	1:500 - 1:2000
IHC	1:50 - 1:200
IF/ICC	1:50 - 1:100

Immunogen Information

Gene ID

9619

Swiss Prot

P45844

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 6-120 of human ABCG1 (NP_004906.3).

Synonyms

ABC8;WHITE1;ABCG1

Contact

 | www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of extracts of various cell lines, using ABCG1 Polyclonal Antibody (A4328) at 1:1000 dilution.


Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (A5014) at 1:10000 dilution.

Lysates/proteins: 25ug per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.



Immunohistochemistry of paraffin-embedded human colon carcinoma using ABCG1 Rabbit pAb (A4328) at dilution of 1:100 (40x lens).



Immunofluorescence analysis of Rat lung using ABCG1 antibody (A4328) at dilution of 1:100. Blue: DAPI for nuclear staining.