Leader in Biomolecular Solutions for Life Science



Catalog No.: A7798 1 Publications



## **Basic Information**

**Observed MW** 69kDa

Calculated MW 68kDa

**Category** Primary antibody

Applications WB,IF/ICC

Cross-Reactivity Human, Mouse

## Background

This gene encodes a member of the acyl-CoA dehydrogenase family. Members of this family of proteins localize to the mitochondria and catalyze the rate-limiting step in the beta-oxidation of fatty acyl-CoA. The encoded protein is specifically active toward palmitoyl-CoA and long-chain unsaturated substrates. Mutations in this gene cause acyl-CoA dehydrogenase family member type 9 deficiency. Alternate splicing results in multiple transcript variants.

## **Immunogen Information**

 WB
 1:500 - 1:2000

 IF/ICC
 1:50 - 1:100

**Recommended Dilutions** 

**Gene ID** 28976 Swiss Prot Q9H845

#### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-270 of human ACAD9 (NP\_054768.2).

### Synonyms

ACAD9;NPD002

# Product Information

www.abclonal.com


**lsotype** IgG Purification Affinity purification

#### Storage

Source

Rabbit

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.



Western blot analysis of extracts of various cell lines, using ACAD9 antibody (A7798) at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25ug per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.