

Affinity Gel-conjugated Mouse anti DDDDK-Tag mAb

Catalog No.: AE061

Basic Information

Observed MW

Refer to Figures

Calculated MW

Category

Tag antibody

Applications

IP

Cross-Reactivity

CloneNo number

AMC0519

Background

FLAG-tag, or FLAG octapeptide, or FLAG epitope, is a polypeptide protein tag that can be added to a protein using recombinant DNA technology, having the sequence motif DYKDDDDK. It has been used for studying proteins in living cells and for protein purification by affinity chromatography. It has been used to separate recombinant, overexpressed protein from wild-type protein expressed by the host organism. It can also be used in the isolation of protein complexes with multiple subunits, because its mild purification procedure tends not to disrupt such complexes. It has been used to obtain proteins of sufficient purity and quality to carry out 3D structure determination by x-ray crystallography. A FLAG-tag can be used in many different assays that require recognition by an antibody. If there is no antibody against a given protein, adding a FLAG-tag to a protein allows the protein to be studied with an antibody against the FLAG sequence. Examples are cellular localization studies by immunofluorescence or detection by SDS PAGE protein electrophoresis and Western blotting.

Recommended Dilutions

Binding Cap 1.1 mg Flag protein/mL

Immunogen Information

Gene ID

Swiss Prot

Immunogen

A synthetic peptide corresponding to DDDDK tag.

Synonyms

DDDDK;DDDDK tag;DDDDK-tag

Contact

 | www.abclonal.com

Product Information

Source

Mouse

Isotype

IgG2b,Kappa

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Validation Data



Immunoprecipitation analysis of 300ug extract cell lysate from 293T cells transfected with GSK3B expression vector containing a DDDDK-tag with 30uL Affinity Gel-conjugated Mouse anti DDDDK-Tag mAb antibody (AE061).Western blot was performed from the immunoprecipitate using Rabbit anti DDDDK-Tag antibody at 1:10000 dilution.



Immunoprecipitation analysis of 300ug extract cell lysate from 293T cells transfected with IRF1 expression vector containing a DDDDK-tag with 30uL Affinity Gel-conjugated Mouse anti DDDDK-Tag mAb antibody (AE061).Western blot was performed from the immunoprecipitate using Rabbit anti DDDDK-Tag antibody at 1:10000 dilution.