

## Recombinant Human Tie2/CD202b Protein (His Tag)(Active)

Catalog No. PKSH031472

### Description

<b>Synonyms</b>	CD202B;TIE-2;TIE2;VMCM;VMCM1
<b>Species</b>	Human
<b>Expression_host</b>	HEK293 Cells
<b>Sequence</b>	Met1-Lys745
<b>Accession</b>	NP_000450.2
<b>Mol_Mass</b>	82 kDa
<b>AP_Mol_Mass</b>	95-105 kDa
<b>Tag</b>	C-His
<b>Bio_activity</b>	Measured by its binding ability in a functional ELISA. Immobilized recombinant human Tie2 at 2 µg/ml (100 µl/well) can bind human Angiopoietin-2 at a linear range of 1.28-160 ng/ml.

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Background

TEK, or TIE-2, is an endothelial cell-specific receptor tyrosine kinase (RTK) that is known as a functioning molecule of vascular endothelial cells. TEK comprises a subfamily of RTK with TIE, and these two receptors play critical roles in vascular maturation, maintenance of integrity and remodeling. Targeted mutagenesis of both Tek and its agonistic ligand, Angiopoietin-1, result in embryonic lethality, demonstrating that the signal transduction pathways mediated by this receptor are crucial for normal embryonic development. TEK signaling is indispensable for the development of the embryonic vasculature and suggests that TEK signaling may also be required for the development of the tumor vasculature.

## SDS-PAGE

