

Recombinant Human CD14 Protein (aa 1-344, His Tag)

Catalog No. PKSH031892

Description

| | |
|------------------------|--|
| Synonyms | Monocyte Differentiation Antigen CD14; Myeloid Cell-Specific Leucine-Rich Glycoprotein; CD14 |
| Species | Human |
| Expression_host | HEK293 Cells |
| Sequence | Met1-Met344 |
| Accession | NP_000582.1 |
| Mol_Mass | 36.5 kDa |
| AP_Mol_Mass | 45 kDa |
| Tag | C-His |

Properties

| | |
|-----------------------|---|
| Purity | > 97 % as determined by reducing SDS-PAGE. |
| Endotoxin | < 1.0 EU per µg as determined by the LAL method. |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. |
| Formulation | Lyophilized from sterile PBS, pH 7.4 |
| Reconstitution | Please refer to the printed manual for detailed information. |

Background

The cluster of differentiation (CD) system is commonly used as cell markers in immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules which associating with the immune function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. Cluster of differentiation 14 (CD14) is a member of the CD system. It takes its name from its inclusion in the CD molecule surface marker proteins. CD14 exists in two forms: a form anchored into the membrane or a soluble form. CD14 was found expressed in macrophages, neutrophil granulocyte and dendritic cells. The major function is serve as a co-receptor (along with TLR4 and MD-2) for the bacterial lipopolysaccharide (LPS) and other pathogen-associated molecular patterns.

SDS-PAGE

