

## Recombinant Cynomolgus CD3d/CD3 delta Protein (Fc Tag)

Catalog No. PKSQ050020

### Description

<b>Synonyms</b>	T-cell surface glycoprotein CD3 delta chain; T-cell receptor T3 delta chain; CD3d; CD3D
<b>Species</b>	Cynomolgus
<b>Expression_host</b>	Human Cells
<b>Sequence</b>	Phe22-Ala105
<b>Accession</b>	Q95LI8
<b>Mol_Mass</b>	36.7 kDa
<b>AP_Mol_Mass</b>	35 kDa
<b>Tag</b>	C-Fc

### Properties

<b>Purity</b>	> 90 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of PBS,pH 7.4.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Background

T-cell surface glycoprotein CD3 delta chain (CD3D) is a single-pass type I membrane protein. CD3D, together with CD3-gamma, CD3-epsilon and CD3-zeta, and the T-cell receptor alpha/beta and gamma/delta heterodimers, forms the T cell receptor-CD3 complex. CD3 chains are present as CD3gammaepsilon, deltaepsilon, and zetaeta dimers in the receptor complex and play critical roles in the antigen receptor assembly, transport to the cell surface, and the receptor-mediated signal transduction. T cell receptor-CD3 complex plays an important role in coupling antigen recognition to several intracellular signal-transduction pathways. This complex is critical for T-cell development and function, and represents one of the most complex transmembrane receptors. The T cell receptor-CD3 complex is unique in having ten cytoplasmic immunoreceptor tyrosine-based activation motifs(ITAMs). CD3D contains 1 ITAM domain and has been shown to interact with CD8A.

## SDS-PAGE

