

## Recombinant Cynomolgus 4-1BB/TNFRSF9/CD137 Protein (Fc Tag)

Catalog No. PKSQ050051

•	4.0
Descri	ntian
DUSCII	

Synonyms CD137; ILA; TNFRSF9; 4-1BB ligand receptor; CDw137; T-cell antigen 4-1BB

homolog; T-cell antigen ILA

SpeciesCynomolgusExpression\_hostHuman CellsSequenceLeu24-Gln186AccessionA9YYE7Mol\_Mass44.4 kDaAP\_Mol\_Mass60 kDa

## **Properties**

Tag

**Purity** > 95 % as determined by reducing SDS-PAGE.

C-Fc

**Endotoxin** < 1.0 EU per μg as determined by the LAL method.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

FormulationLyophilized from a 0.2 μm filtered solution of PBS,pH 7.4.ReconstitutionPlease refer to the printed manual for detailed information.

## Background

Tumor necrosis factor receptor superfamily member 9(TNFRSF9), also known as CD137 and 4-1BB, is an inducible T cell surface protein belonging to the tumor necrosis factor receptor superfamily. It is a single-pass type I membrane protein which contains 4 TNFR-Cys repeats. The human and mouse proteins share 60% amino acid sequence identity. CD137 is expressed by mesenchymal cells, including endothelial cells, chondrocytes, and cells of the central nervous system. CD137 is also broadly expressed by cells of the human immune system, is broadly expressed by cells of the human immune system, including activated CD8+ and CD4+ T cells, activated natural killer (NK) cells, follicular dendritic cells (FDCs) and monocytes. CD137 has diverse roles in the immune response, the one key function is to promote the survival of both T cells and dendritic cells by binding the cognate ligand CD137L (4-1BBL).



## SDS-PAGE

