

Recombinant Mouse CXCL2/MIP-2 Protein

Catalog No. PKSM040996

Description

Synonyms	MIP-2; chemokine ligand 2; C-X-C motif chemokine 2; GRO beta; GRO2; GROB; Gro-beta; Growth-regulated protein beta; Macrophage Inflammatory Protein-2-alpha; melanoma growth stimulatory activity beta; cxcl2; MGSA-b; MGSA-beta; MIP2A; MIP2-alpha; SCYB2.
Species	Mouse
Expression_host	E.coli
Sequence	Ala28-Asn100
Accession	P10889
Mol_Mass	7.9 kDa
AP_Mol_Mass	10 kDa
Tag	No tag

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM Tris, 150mM NaCl, pH8.0.
Reconstitution	Please refer to the printed manual for detailed information.

Background

C-X-C motif chemokine 2 (CXCL2, MIP-2) belongs to the intercrine alpha (chemokine Cx) family. It was originally identified as a heparin-binding protein secreted from a murine macrophage cell line in response to endotoxin stimulation. The expression of mouse MIP-2 is stimulated by endotoxin. The mouse MIP-2 shares approximately 63% aa sequence identity with murine KC, another mouse alpha chemokine, which is induced by PDGF. It has been suggested that mouse KC and MIP-2 are the homologs of the human GROs and rat CINC. Chemotactic for human polymorphonuclear leukocytes but does not induce chemokinesis or an oxidative burst. The expression of MIP-2 was found to be associated with neutrophil influx in pulmonary inflammation and glomerulonephritis, suggesting that MIP-2 may contribute to the pathogenesis of inflammatory diseases.

SDS-PAGE

