

## Recombinant Mouse CD16-2/FCGR4 Protein (His&AVI Tag), Biotinylated(Active)

Catalog No. PKSM040891

### Description

<b>Synonyms</b>	4833442P21Rik Protein, Mouse;CD16-2 Protein, Mouse;FcgammaRIV Protein, Mouse;Fcgr3a Protein, Mouse;FcgrIV Protein, Mouse;Fcrl3 Protein, Mouse
<b>Species</b>	Mouse
<b>Expression_host</b>	HEK293 Cells
<b>Sequence</b>	Met1-Gln203
<b>Accession</b>	NP_653142.2
<b>Mol_Mass</b>	24.2 kDa
<b>Tag</b>	C-His-Avi, Biotinylated
<b>Bio_activity</b>	1. Measured by its ability to bind human IgG1 (Fc) in a functional ELISA.2. Labeling ratio of biotin to protein: 0.6

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Background

Fcgr4, also known as CD16-2, is one of the receptors for Fc region of IgG which involve in immune responses. Fcgr4 mainly functions in cellular response to lipopolysaccharide, NK T cell proliferation, regulation of sensory perception of pain, wound healing etc. Three groups are included for Fc γ receptors (FcR), and they are Fc γ RI (CD64), Fc γ RII (CD32), and Fc γ RIII (CD16). Among these, CD64 possess high affinity even for monomeric IgG, while CD32 and CD16 display a relative lower affinity for IgG. Genes encode these receptors are diverse differing by species and cell types. The aggregation of FcR having immunoreceptor tyrosine-based activation motifs (ITAMs) activates sequentially src family tyrosine kinases and syk family tyrosine kinases that connect transduced signals to common activation pathways shared with other receptors. FcR with ITAMs elicit cell activation, endocytosis, and phagocytosis. Fcgr4 belongs to Fc γ RIII (CD16) group.

## SDS-PAGE

