


Tel:240-252-7368(USA) Fax: 240-252-7376(USA) techsupport@elabscience.com Website: www.elabscience.com

AADAC Polyclonal Antibody


Catalog No.E-AB-14453ReactivityHStorageStore at -20°C. Avoid freeze / thaw cycles.HostRabbitApplicationsWB,IHC,ELISAIsotypeIgG

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Images



Western Blot analysis of Human fetal liver and liver cancer tissue using AADAC Polyclonal Antibody at dilution of 1:550



Immunohistochemistry of paraffinembedded Human colon cancer using AADAC Polyclonal Antibody at dilution of 1:30



Immunohistochemistry of paraffinembedded Human liver cancer using AADAC Polyclonal Antibody at dilution of 1:30

Immunogen Information

Immunogen Recombinant protein of human AADAC

Gene Accession BC032309 Swissprot P22760 Synonyms DAC,CES5A1

Product Information

Calculated MW 46kDa

Buffer PBS with 0.05% sodium azide and 50% glycerol,

PH7.4

Purify Affinity purification

Dilution WB 1:500-1:2000, IHC 1:25-1:100

Background

Microsomal arylacetamide deacetylase competes against the activity of cytosolic arylamine N-acetyltransferase, which catalyzes one of the initial biotransformation pathways for arylamine and heterocyclic amine carcinogens. Arylacetamide deacetylation is an important enzyme activity in the metabolic activation of arylamine substrates to ultimate carcinogens. Displays major serine hydrolase activity in liver microsomes. Hydrolyzes also flutamide, which is an antiandrogen drug used for the treatment of prostate cancer that occasionally causes severe hepatotoxicity. Displays cellular triglyceride lipase activity in liver. Increases intracellular fatty acids derived from hydrolysis of newly formed triglyceride stores.

For Research Use Only

Thank you for your recent purchase

If you would like to learn more about antibodies, please visit www.elabscience.com.

Focus on your research Service for life science