

Active Recombinant Human IL-17A/CTLA-8 Protein

Catalog No.: RP00212 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Human	3605	Q16552

Tags

C-His

Synonyms

CTLA8; IL17; IL17A

Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE.

Endotoxin

< 0.1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

Reconstitution

Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water.

Background

IL17, also known as IL17a and CTLA8, is a 15-20 kDa glycosylated cytokine which belongs to the IL-17 family. The IL-17 family of cytokines includes six members, IL-17/IL-17A, IL-17B, IL-17C, IL-17D, IL-17E/IL-25, and IL-17F, which are produced by multiple cell types. IL17A promotes protective mucosal and epidermal inflammation in response to microbial infection. It induces chemokine production, neutrophil influx, and the production of antibacterial peptides. IL17A additionally enhances the production of inflammatory mediators by rheumatoid synovial fibroblasts and contributes to TNF alpha induced shock. In contrast, it can protect against the progression of colitis by limiting chronic inflammation. IL17A encourages the formation of autoreactive germinal centers and exacerbates the onset and progression of experimental models of autoimmunity.

Basic Information

Description

Active Recombinant Human IL-17A/CTLA-8 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Ile20-Ala155) of human IL-17A (Accession #NP_002181.1) fused with a 6xHis tag at the C-terminus.

Bio-Activity

1. Measured by its ability to induce IL6 secretion in Hela cells. The ED50 for this effect is 1-6 ng/mL.
2. Measured by its binding ability in a functional ELISA. Immobilized Recombinant human IL17RA at 2 μg/mL (100 μL/well) can bind Recombinant human IL-17A, the EC50 of human IL-17A is 32.5-130 ng/mL.


Storage

Store the lyophilized protein at -20°C to -80 °C for long term. After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week. Avoid repeated freeze/thaw cycles.


Contact

 | www.abclonal.com


Validation Data



Recombinant Human IL-17A Protein was determined by SDS-PAGE with Coomassie Blue, showing bands at 18-25 kDa.



Recombinant Human IL-17A induces IL6 secretion in Hela cells. The ED_{50} for this effect is 1-6 ng/mL.



Immobilized Recombinant human IL17RA , His Tag at 2 μ g/mL (100 μ L/well) can bind Recombinant human IL-17A, The EC_{50} of human IL-17A is 32.5-130 ng/mL.