Mouse Negative Control





www.biosb.com

Intended Use

For In Vitro Diagnostic Use.

Summary and Explanation

The purpose of the Mouse Negative Control is to demonstrate whether the IHC test is specific. Usually, "negative control" is a term applied to a separate slide with patient's test sample on which the Mouse Negative Control is applied.

Presentation

The Mouse Negative Control is provided in liquid form ready-to-use with Tris Buffered Saline, pH 7.6, 1% bovine serum albumin and preserved with Sodium Azide

Catalog No.	Concentration	Volume
BSB 0040A	Ready-to-use	3 mL
BSB 0040B	Ready-to-use	6 mL
BSB 0040C	Ready-to-use	15 mL

Storage Store at 2-8°C

Stability

This product is stable up to the expiration date on the product label. Do not use after expiration date listed on package label. Temperature fluctuations should be avoided. Store appropriately when not in use.

Precautions

- 1 For professional users only. Results should be interpreted by a medical professional.
- 2. This product contains <0.1% sodium azide (NaN₃) as a preservative. Ensure proper handling procedures are used with this reagent.
- 3. Always wear personal protective equipment such as laboratory coat, goggles and gloves when handling reagents.
- 4. Dispose of unused solution according to local and federal regulations.
- 5. Do not ingest reagent. If reagent is ingested, seek medical advice immediately.
- $6.\ Avoid\ contact\ with\ eyes.\ If\ contact\ occurs,\ flush\ with\ large\ quantities\ of\ water.$
- 7. For complete recommendations for handling biological specimens please refer to the CDC document, "Guidelines for Safe Work Practices in Human and Animal Medical Diagnostic Laboratories" (1).

Preparation of Working Solution

The Mouse Negative Control is a ready-to-use working solution and requires no further preparation.

Recommended Protocol

In order to serve as a true negative control, this mouse antibody should be used under the identical conditions and at the same concentration as other purified mouse monoclonal antibodies.

Mouse negative control serum is applied in place of a mouse anti-human primary antibody onto tissue that has been designated as the corresponding negative control. The section under test and the corresponding negative control should be serial sections off the same paraffin-embedded block.

Note:

Both the tissue under test and the negative control tissue must be unmasked using the same unmasking reagent and protocol such as Heat Induced Epitope Retrieval (HIER).

Abbreviated Immunohistochemical Protocol

Step	ImmunoDetector AP/HRP	PolyDetector AP/HRP	PolyDetector Plus HRP		
Peroxidase/AP Blocker	5 min.	5 min.	5 min		
Primary Antibody	30-60 min.	30-60 min.	30-60 min. 15 min. 15 min.		
1st Step Detection	10 min.	30-45 min.			
2nd Step Detection	10 min.	Not Applicable			
Substrate-Chromogen	5-10 min.	5-10 min.	5-10 min.		
Counterstain / Coverslip	Varies	Varies	Varies		

Product Limitations

Due to inherent variability present in immunohistochemical procedures (including fixation time of tissues, dilution factor of antibody, retrieval method utilized and incubation time), optimal performance should be established through the use of positive and negative controls. Results should be interpreted by a qualified medical professional.

References

1. U.S. Department of Health and Human Services: Centers for Disease Control and Prevention. Guidelines for Safe Work Practices in Human and Animal Medical Diagnostic Laboratories. Supplement / Vol. 61, January 6, 2012.

Symbol Key / Légende des symboles/Erläuterung der Symbole

ر	Symbol key / Legende des symboles/Enduterding der Symbole									
		n esc	Storage Temperature		Manufacturer		Catalog Number			
			Limites de température		Fabricant	REF	Référence du catalogue			
		2.0	Zulässiger Temperaturbereich		Hersteller		Bestellnummer			
	In Vitro Diagnostic Medical Device	~	Read Instructions for Use		Expiration Date	LOT	Lot Number			
	Dispositif médical de diagnostic in vitro	 _ -	Consulter les instructions d'utilisation	><	Utiliser jusque	101	Code du lot			
	In-Vitro-Diagnostikum		Gebrauchsanweisung beachten		Verwendbar bis		Chargenbezeichnung			